15

20

WHAT IS CLAIMED IS:

- 1. A fungicidal composition comprising a synergistic fungicidally effective amount of a compound A which is (i) methyl (E)-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate or (ii) N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxymethyl)phenyl]acetamide, and a compound B which is iprodione, the A/B ratio by weight being between 0.02 and 5.
- 2. A fungicidal composition according to Claim 1, wherein compound A is methyl (E)-methoxyimino [α -(o-tolyloxy)-o-tolyl]acetate.
- 3. A fungicidal composition according to Claim 1, wherein compound A is N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxymethyl)phenyl]acetamide.
- 4. A fungicidal composition according to Claim 1, wherein the A/B ratio by weight is between 0.0625 and 1.33.

5. A fungicidal composition according to Claim 1, wherein the A/B ratio by weight is between 0.05 and 5.

- A fungicidal composition according to Claim 1,
 wherein the A/B ratio by weight is between 0.17 and
 1.33.
 - 7. A fungicidal composition according to Claim 1, wherein the A/B ratio by weight is between 0.02 and 2.
 - 8. A fungicidal composition according to Claim 1, wherein the A/B ratio by weight is between 0.0625 and 0.25.

10

15

20

25

- 9. A fungicidal composition according to Claim 1, further comprising at least one member selected from the group consisting of an agriculturally acceptable solid or liquid vehicle and an agriculturally acceptable surface-active agent.
- 10. A fungicidal composition according to Claim 9, comprising from 0.05% to 95% by weight of compounds A and B.
- 11. A process for the curative or preventative control of phytopathogenic fungi in plants, said process comprising applying to plants or to the locus in which they grow a synergistic fungicidally effective non-phytotoxic amount of a compound A which is (i) methyl (E)-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate or (ii) N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxymethyl)phenyl]acetamide, and a compound B which is iprodione, the A/B ratio by weight being between 0.02 and 5.
- 12. A process according to Claim 11, wherein compound A is methyl (E)-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate.
- 13. A process according to Claim 11, wherein compound A is N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxymethyl)phenyl]acetamide.
- 30 14. A process according to Claim 11, wherein the A/B ratio by weight is between 0.0625 and 1.33.
 - 15. A process according to Claim 11, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to crops or lawns to control fungi

10

15

20

25

30

which are phytopathogenic towards crops or lawns, respectively.

- 16. A process according to Claim 15, wherein compound A is methyl-(E)-methoxyimino[α -(o-tolyloxy)-o-tolyl]acetate.
 - 17. A process according to Claim 15, wherein compound A is N-methyl-(E)-methoxyimino[2-(2,5-dimethylphenoxymethyl)phenyl]acetamide.
 - 18. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to the aerial parts of the plants.
 - 19. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to crops and wherein the A/B ratio by weight is between 0.05 and 5.
 - 20. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to crops and wherein the A/B ratio by weight is between 0.17 and 1.33.
 - 21. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to crops at a dose rate of between 150 and 1500 g/ha.
 - 22. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to crops at a dose rate of between 400 and 1000 g/ha.

- 23. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to lawns and wherein the A/B ratio by weight is between 0.02 and 2.
- 24. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to lawns and wherein the A/B ratio by weight is between 0.0625 and 0.25.
- $25.\,$ A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to lawns at a dose rate of between 1100 and 7000 g/ha.
- 26. A process according to Claim 15, wherein the synergistic fungicidally effective non-phytotoxic amount of A and B is applied to lawns at a dose rate of between 2250 and 5000 g/ha.